

NOTES FOR SEPTEMBER 17: JACOBI-TRUDI IDENTITY

CHRIS FRASER

Our goal today is to express s_λ in terms of $\{h_\mu : \mu \vdash n\}$. We do so via the following:

Theorem 1. (*Jacobi-Trudi Identity*): For λ satisfying $\ell(\lambda) \leq n$, we have

$$s_\lambda = \det((h_{\lambda_i - i + j})_{i,j \in [n]}).$$

Here, $h_0 = 1, h_k = 0$ for $k < 0$.

We will refer to this henceforth as JT.


Example 1. JT says that

$$\begin{aligned} s_{321} &= \det \begin{pmatrix} h_3 & h_4 & h_5 \\ h_1 & h_2 & h_3 \\ 0 & h_0 & h_1 \end{pmatrix} \\ &= h_{321} - h_{411} - h_{330} + h_{510}. \end{aligned}$$

The way to “remember” the statement of the theorem is to note that the indices of the the h_i ’s along the diagonal are the parts of λ , and that the indices increase by 1 as they move right. For example: we have h_3, h_2, h_1 along the diagonal in the preceding example.

Proof. Fix some $N \gg 0$. Let the “top vertices” denote the n integer lattice points with coordinates $(0, N), (1, N), \dots, (n-1, N)$. Let the “bottom vertices” denote the n points with coordinates $(\lambda_n, 1), (\lambda_{n-1} + 1, 1), (\lambda_{n-2} + 2, 1), \dots, (\lambda_1 + (n-1), 1)$. Read these bottom points from left to right, so that $(\lambda_n, 1)$ is the first point, and $(\lambda_1 + n - 1, 1)$ is the last point. In the figure below, we have taken $n = 3$ and $N = 5$. The top vertices and bottom vertices are the labeled lattice points. λ is the partition $\lambda = (532)$.

We want to interpret both the LHS and RHS of JT as weighted sums of collections of n lattice paths joining the “top vertices” to the “bottom vertices”. Namely, we are considering **lattice n -paths**, $L = (L_1, \dots, L_n)$, joining the top points to the bottom points. Each L_i is a lattice path joining $(i-1, N)$ to the $\sigma(i)^{th}$ special point on the bottom, for some $\sigma \in \mathfrak{S}_n$. Each L_i is only allowed to move down or right. The figure below shows a lattice 3-path with $\sigma = \text{id}$.



We weight horizontal edges $(k, j) \rightarrow (k+1, j)$ in the integer lattice with the monomial x_j , and give all vertical edges have weight 1. Then the **weight** of a given path L_j is the product of its weights, which is

$$w(L_j) := \prod_{\text{horizontal steps of } L_j \text{ at height } i} x_i.$$

The *weight* of a lattice n -path, $L = (L_1, \dots, L_n)$ is just the product of the weights of its paths:

$$w(L) := \prod_i w(L_i).$$

The weights of $L(1), L(2), L(3)$ in the diagram above are x_2^2, x_3^3 and $x_4^3 x_3^2$ respectively. $w(L)$ is consequently $x_2^2 x_3^5 x_4^3$.

Now let's return to the proof.

Claim 1. For a fixed $\sigma \in \mathfrak{S}_n$:

a)

$$\sum_{L \text{ joins top to bottom according to } \sigma} w(L) = \prod_{i=1}^n h_{\lambda_{\sigma(i)} - \sigma(i) + i}.$$

b) The RHS of JT is

$$\sum_{L=(L_1, \dots, L_n)} \epsilon_{\sigma(L)} w(L).$$

To see a): First, we note that

$$\sum_{L_j \text{ has horizontal distance } k} w(L_j) = h_k.$$

Indeed, by definition,

$$h_k(x_1, \dots, x_n) = \sum_{i_1 \leq i_2 \leq i_n} x_1^{i_1} \cdots x_n^{i_n}$$

and we associate to the sequence $(i_1 \leq i_2 \leq i_n)$ the path that takes i_n steps right, then moves down 1, then takes i_{n-1} steps right, then moves down 1, etc. In the end, we move k steps right. The resulting path has weight $x_1^{i_1} \cdots x_n^{i_n}$.

Once you know this, observe that paths of type σ have L_i moving $\lambda_{\sigma(i)} - \sigma(i) + i$ steps right, from which we obtain a), since the weight of the collection (L_1, \dots, L_n) is just the product of their individual weights.

To see b): Follows from the definition of the determinant as a sum over permutations, using part a).

Claim 2. $s_\lambda = \sum_L \epsilon_{\sigma(L)} w(L) = \text{RHS of JT, by Claim 1.}$

To do this, we note that n -lattice paths in which some L_i intersects another L_j cancel out in pairs. Let's do this carefully: pick the topmost and leftmost point of intersection of L_i and L_j . We define a new lattice n -path \tilde{L} as follows. Every path in L besides paths L_i and L_j becomes a path in \tilde{L} . We must specify two more paths to complete our description of \tilde{L} . In the first path, we follow L_i until the point of intersection, after which we follow L_j . In the second path, we follow L_j until the intersection, after which we follow L_i . Then $w(L) = w(\tilde{L})$ since all of the same edges are taken. However the the permutations associated to the two paths differs by the transposition (ij) , so that their signs are opposite, and the contributions of L and \tilde{L} to the above weighted sum cancel out.

The remaining paths are the *nonintersecting* lattice n -paths. Note that any such lattice n -path corresponds to the identity permutation ("topologically" obvious: any nontrivial permutation would have an inversion, and the two paths corresponding to this inversion would have to cross). The drawing of a 3-lattice path we gave above is an example of such a nonintersecting path.

We claim nonintersecting lattice n -paths of shape λ are in weight-preserving bijection with *reverse SSYT* of shape λ . A *reverse SSYT* is a filling of λ that is weakly decreasing in rows and strictly decreasing in columns. We fill the first row of λ by reading off the horizontal edges of L_n in decreasing order. The second row of λ is filled with the horizontal edges of L_{n-1} ,

and so on. For example, the 3-lattice path we drew above corresponds to the Reverse SSYT

4	4	4	3	3
3	3	3		
2	2			

Clearly, the filling we have prescribed is decreasing along rows. That it is strictly decreasing in columns amounts to the assertion that the paths do not self-intersect. This process is reversible, so we have the claimed bijection.

By the preceding, we will have proven claim 2, and consequently the JT theorem, after the following exercise:

Exercise: $s_\lambda = \sum_{T \in \text{ReverseSSYT}(\lambda)} x^T$. (Hint, fill each box in a reverse SSYT by n minus what is currently in the box).

□

Remark: Jacobi-Trudi generalizes to skew shapes:

$$s_{\lambda/\mu} = \det((h_{\lambda_i - \mu_i - i + j})_{i,j \in [n]}).$$

Here, the definition of $s_{\lambda/\mu}$ is the same as the combinatorial definition of the s_λ : it is the content-generating function for SSYT of shape λ/μ . The proof of this identity is the same as above, where one uses the parts of μ to define the values of the “top points”, just as we used the value of λ to define the “bottom points” in our proof above.

Remark: I believe we have not yet discussed that the involution ω sends $s_\lambda \mapsto s_{\lambda'}$. Once we see this, we will obtain as a corollary

$$s_\lambda = \det((e_{\lambda'_i} - i + j)_{i,j \in [n]}),$$

which follows by applying ω to both sides of JT.

David adds: Actually, my plan was to have Jonah also mention the dual identity stated here, point out that it can be proved the same way, and deduce from that that ω sends s_λ to $s_{\lambda'}$.